

Flux Monitoring Using the Toruń 32-m Radio Telescope

J. Kijak

Institute of Astronomy, University of Zielona Góra, Poland

Y. Gupta

NCRA, TIFR, Pune University Campus, Pune, India

W. Lewandowski, G. Hrynek

TCfA, Nicolaus Copernicus University, Toruń, Poland

K. Krzeszowski

Institute of Astronomy, University of Zielona Góra, Poland

Abstract. We present flux density measurements for PSR B0329+54 at 4.8 GHz obtained during one year.

1. Observations and Results

We have observed PSR B0329+54 during one year using the Toruń 32-m radio telescope. Ten epochs of flux monitoring at 4.8 GHz distributed over one year are reported here. Typical observations lasted about 24 hours in 3 minutes scans, but two epochs (Fig. 1) had longer observation times (5 and 2 days, respectively). The total intensity signal was obtained by summing the dual-circular polarization signals, and dedispersing over 64 channels covering the 192 MHz bandwidth. In order to convert to flux density we carried out regular calibration measurements using the injected signal of a noise diode, which was compared to the flux density of known continuum calibrators. The observations were interrupted regularly for calibration procedure and sometimes due to technical problems. We have obtained a combined time series of calibrated flux measurements for each epoch. Typical errors of flux measurements are about 10%. The longer epochs of flux density time series are presented in Figure 1.

2. Analysis and Conclusions

For each epoch, the average flux density and modulation index were calculated. The average flux density S_{av} for the one year period is 16 ± 6 mJy, which is consistent with the known spectrum of this pulsar (Maron et al. 2000). The modulation index m varies between 0.7 and 1.3. To estimate the modulation timescale, we computed the structure functions for our data sets. The estimated timescale is 90 minutes, with an uncertainty of about 15 minutes.




Figure 1. Two epochs of flux density time series. Note a significant difference in fluxes between these two epochs separated by about six months.

Summarizing, (1) we conclude from our preliminary results that the data quality is good enough for detailed studies of ISS of PSR B0329+54 at 5 GHz, (2) our analysis shows that the modulation index m is close to unity, consistent with Malofeev et al. (1996) who reported $m \geq 0.5$. However, our time scale of 90 min is significantly larger than the 21 min given by Malofeev et al. (1996). Our result is much closer to the time scale of 80 min estimated by Kramer et al. (2003), (3) further analysis of multi-epoch data spread over more than one year should result in better understanding of the scattering geometry for PSR B0329+54.

Acknowledgments. This work was supported in part by the Polish State Committee for Scientific Research Grant 2 P03D 008 19. JK acknowledges partial support from the IAU.

References

- Kramer, M., Karastergiou, A., Gupta, Y., Johnston, S., Bhat, N. D. R., & Lyne, A. G. 2003, *A&A*, 407, 655
 Malofeev, V. M., Shishov, V., Sieber, W., Jessner, A., Kramer, M., & Wielebinski, R. 1996, *A&A*, 308, 180
 Maron, O., Kijak, J., Kramer, M., & Wielebinski, R. 2000, *A&AS*, 147, 195